

Meta-analyses published between 2000 and 2025 on alcohol consumption and all-cause mortality

Author (Year)	Total number of studies or cohorts	Same or lower risk	J-shaped risk curve	Former drinking is separated from lifetime abstinence
Stockwell (2024) ¹	107 studies published between 1950s and 31 July 2021	✓	✗	✓
Zhao (2023) ^{2,3}	107 studies published between 1950s and 31 July 2021	✓	✗	✓
Stockwell (2016) ²	87 studies published between 1950s and 31 December 2014	✓	✗	✓
Fillmore (2006)	54 studies published between 1950s and mid-2004	✓	✗	✓
Di Castelnuovo (2023) ¹	16 cohorts recruited as early as 1983 with up to 30 years of follow-up	✓	✓	✓
Di Castelnuovo (2022)	16 cohorts recruited as early as 1983 with up to 30 years of follow-up	✓	✓	✓
Hu et (2022) ⁴	3 cohorts recruited as early as 1998 with an average follow-up of 7.5 years	✓	✓	✗
Degerud (2020)	2 cohorts recruited between 1994 and 2002 with an average follow-up of 16.7 years	✓	✓	✗
Colpani (2018) ⁵	59 studies published up to 29 February 2016	✓	✓	✗
Li (2018)	2 cohorts recruited as early as 1980 with an average follow-up of 34 years	✓	✓	✗
Wood (2018)	83 studies published up to 1 March 2017	✓	✓	✓
Perreault (2017) ^{1,3}	8 cohorts recruited as early as 1994 with an average follow-up of 9.7 years	✓	✓	✓
Soedamah-Muthu (2013)	4 cohorts recruited as early as 1997 with an average follow-up of 9.7 years	✓	✓	✗
Bobak (2016) ³	4 cohorts recruited as early as 2002 with an average follow-up of 6.9 years	✓	✓	✗
Zheng (2015) ³	23 studies published through June 2014	✓	✓	✗

Continued on next page

Continued

Ferrari (2014) ^{1,3}	23 cohorts recruited as early as 1992 with an average follow-up of 12.6 years	✓	✓	✓
Sluik (2014)	23 cohorts recruited as early as 1992 with an average follow-up of 9.9 years	✓	✓	✗
Jayasekara (2014) ⁴	9 studies published through August 2012	✓	✓	✓
Inoue (2012) ³	6 cohorts recruited as early as 1988 with an average follow-up of 12.4 years	✓	✓	✓
Ronksley (2011)	31 studies published from 1950 through September 2009	✓	✓	✗
Gronbaek (2004) ¹	4 cohorts recruited as early as 1964 with up to 31 years of follow-up	✓	✓	✗
Gronbaek (2000)	8 cohorts recruited as early as 1964 with up to 31 years of follow-up	✓	✓	✗
Rehm (2001) ³	59 studies published between 1975 and 2001	✓	✓	✗

KEY:  YES  NO**INCLUSION CRITERIA**

This analysis included studies that met the following criteria:

- Study design: meta-analysis or pooled cohort
- Study population: must include a general population, rather than a subpopulation with an existing condition or diagnosis (for example, diabetes or hypertension patients)
- Exposure: must include total alcohol consumption (rather than beverage-specific results only) and provide estimates for different levels of alcohol consumption (rather than a single binary variable or a single dose-response estimate)
- Outcome: all-cause mortality
- Analysis: must contain a comparison of alcohol consumption to no alcohol consumption
- Publication date: 2000 to March 2025

Note: Many of the meta-analyses included in this table have similar inclusion criteria. Therefore, they include overlapping sets of individual research studies.

COLUMN NOTES

Total number of studies or cohorts: this total number reflects how many studies met the publication's inclusion criteria. The number of studies that authors used for the analyses or models represented in other columns may differ.

Same or lower risk: studies that found the risk for light-to-moderate alcohol consumption was the same or lower than the risk for no alcohol consumption.

J-shaped risk curve: studies that found a statistically significant lower risk for light-to-moderate alcohol consumption, compared to no alcohol consumption, but not necessarily a statistically significant higher risk for heavier alcohol consumption.

Former drinking is separated from lifetime abstinence: studies that separate former drinking from lifetime abstinence or no alcohol consumption are better able to control for consumers who may have reduced or quit drinking because of an illness, which may bias estimates of risk. Researchers may also test for this potential bias and present results from a combined no alcohol consumption group, if combining the two does not affect the results.

STUDY NOTES

¹ This study draws from the same study population as the ones listed below but may use a sub-sample of the study population, a larger sample of the study population, or a different analysis methodology.

² This study is an update of previously published meta-analyses (Stockwell et al., 2016, and Fillmore et al., 2006), which covers previously included studies, adds more recent studies, and refines coding manuals for data extractions.

³ This study stratified its analysis by age, study population, race, sex, or a combination of those factors. The results of at least one subgroup analyses met the criteria for the same or lower risk compared to no alcohol consumption, a J-shaped risk curve, or separating former drinking from lifetime abstinence as described in the notes.

⁴ This study examines men only.

⁵ This study examines women only.

REFERENCES

- Bobak, M., Malyutina, S., Horvat, P., Pajak, A., Tamosiunas, A., Kubinova, R., et al. (2016). Alcohol, drinking pattern and all-cause, cardiovascular and alcohol related mortality in Eastern Europe. *European Journal of Epidemiology*, 31, 21–30.
- Colpani, V., Baena, C. P., Jaspers, L., van Dijk, G. M., Farajzadegan, Z., Dhana, K., et al. (2018). Lifestyle factors, cardiovascular disease and all-cause mortality in middle-aged and elderly women: A systematic review and meta-analysis. *European Journal of Epidemiology*, 33(9), 831–845.
- Degerud, E., Høiseth, G., Mørland, J., Ariansen, I., Graff-Iversen, S., Ystrom, E., et al. (2020). Association of coincident self-reported mental health problems and alcohol intake with all-cause and cardiovascular disease mortality: A Norwegian pooled population analysis. *PLOS Medicine*, 17(2), e1003030.
- Di Castelnuovo, A., Bonaccio, M., Costanzo, S., McElduff, P., Linneberg, A., Salomaa, V., et al. (2023). Drinking alcohol in moderation is associated with lower rate of all-cause mortality in individuals with higher rather than lower educational level: Findings from the MORGAM project. *European Journal of Epidemiology*, 38(8), 869–881.
- Di Castelnuovo, A., Costanzo, S., Bonaccio, M., McElduff, P., Linneberg, A., Salomaa, V., et al. (2022). Alcohol intake and total mortality in 142,960 individuals from the MORGAM Project: A population-based study. *Addiction*, 117(2), 312–325.
- Ferrari, P., Licaj, I., Muller, D. C., Kragh Andersen, P., Johansson, M., Boeing, H., et al. (2014). Lifetime alcohol use and overall and cause-specific mortality in the European Prospective Investigation into Cancer and nutrition (EPIC) study. *BMJ Open*, 4(7), e005245.
- Fillmore, K. M., Kerr, W. C., Stockwell, T., Chikritzhs, T., & Bostrom, A. (2006). Moderate alcohol use and reduced mortality risk: Systematic error in prospective studies. *Addiction Research & Theory*, 14(2), 101–132.
- Gronbaek, M., Johansen, D., Becker, U., Hein, H. O., Schnohr, P., Jensen, G., et al. (2004). Changes in alcohol intake and mortality: A longitudinal population-based study. *Epidemiology*, 15(2), 222–228.
- Gronbaek, M., Becker, U., Johansen, D., Gottschau, A., Schnohr, P., Hein, H. O., et al. (2000). Type of alcohol consumed and mortality from all causes, coronary heart disease, and cancer. *Annals of Internal Medicine*, 133(6), 411–419.
- Hu, C., Huang, C., Li, J., Liu, F., Huang, K., Liu, Z., et al. (2022). Causal associations of alcohol consumption with cardiovascular diseases and all-cause mortality among Chinese males. *American Journal of Clinical Nutrition*, 116, 771–779.
- Inoue, M., Nagata, C., Tsuji, I., Sugawara, Y., Wakai, K., Tamakoshi, A., et al. (2012). Impact of alcohol intake on total mortality and mortality from major causes in Japan: A pooled analysis of six large-scale cohort studies. *Journal of Epidemiology and Community Health*, 66(5), 448–456.
- Jayasekara, H., English, D. R., Room, R., & MacInnis, R. J. (2014). Alcohol consumption over time and risk of death: A systematic review and meta-analysis. *American Journal of Epidemiology*, 179(9), 1049–1059.
- Li, Y., Pan, A., Wang, D. D., Liu, X., Dhana, K., Franco, O. H., et al. (2018). Impact of healthy lifestyle factors on life expectancies in the US population. *Circulation*, 138, 345–355.
- Perreault, K., Bauman, A., Johnson, N., Britton, A., Rangul, V., & Stamatakis, E. (2017). Does physical activity moderate the association between alcohol drinking and all-cause, cancer and cardiovascular diseases mortality? A pooled analysis of eight British population cohorts. *British Journal of Sports Medicine*, 51(8), 651–657.
- Rehm, J., Gutjahr, E., & Gmel, G. (2001). Alcohol and all-cause mortality: A pooled analysis. *Contemporary Drug Problems*, 28(3), 337–361.
- Ronksley, P. E., Brien, S. E., Turner, B. J., Mukamal, K. J., & Ghali, W. A. (2011). Association of alcohol consumption with selected cardiovascular disease outcomes: A systematic review and meta-analysis. *British Medical Journal*, 342(7795), 479.
- Sluik, D., Boeing, H., Li, K., Kaaks, R., Johnsen, N. F., Tjønneland, A., et al. (2014). Lifestyle factors and mortality risk in individuals with diabetes mellitus: Are the associations different from those in individuals without diabetes? *Diabetologia*, 57(1), 63–72.
- Soedamah-Muthu, S. S., De Neve, M., Shelton, N. J., Tielemans, S. M., & Stamatakis, E. (2013). Joint associations of alcohol consumption and physical activity with all-cause and cardiovascular mortality. *American Journal of Cardiology*, 112(3), 380–386.
- Stockwell, T., Zhao, J., Clay, J., Levesque, C., Sanger, N., Sherk, A., et al. (2024). Why do only some cohort studies find health benefits from low volume alcohol use? A systematic review and meta-analysis of study characteristics that may bias mortality risk estimates. *Journal of Studies on Alcohol and Drugs*, 85(4), 441–452.
- Stockwell, T., Zhao, J., Panwar, S., Roemer, A., Naimi, T., & Chikritzhs, T. (2016). Do “moderate” drinkers have reduced mortality risk? A systematic review and meta-analysis of alcohol consumption and all-cause mortality. *Journal of Studies on Alcohol and Drugs*, 77(2), 185–198.
- Wood, A. M., Kaptoge, S., Butterworth, A. S., Willeit, P., Warnakula, S., Bolton, T., et al. (2018). Risk thresholds for alcohol consumption: Combined analysis of individual-participant data for 599,912 current drinkers in 83 prospective studies. *The Lancet*, 391(10129), 1513–1523.
- Zhao, J., Stockwell, T., Naimi, T., Churchill, S., Clay, J., & Sherk, A. (2023). Association between daily alcohol intake and risk of all-cause mortality: A systematic review and meta-analyses. *JAMA Network Open*, 6(3), e236185.
- Zheng, Y. L., Lian, F., Shi, Q., Zhang, C., Chen, Y. W., Zhou, Y. H., et al. (2015). Alcohol intake and associated risk of major cardiovascular outcomes in women compared with men: A systematic review and meta-analysis of prospective observational studies. *BMC Public Health*, 15(1), 773. Network Open, 6(3), e236185. doi:10.1001/jamanetworkopen.2023.6185